

## ORTHOGONAL LAURENT POLYNOMIALS AND QUADRATURES ON THE UNIT CIRCLE AND THE REAL HALF-LINE\*

RUYMÁN CRUZ-BARROSO AND PABLO GONZÁLEZ-VERA †

**Abstract.** The purpose of this paper is the computation of quadrature formulas based on Laurent polynomials in two particular situations: the Real Half-Line and the Unit Circle. Comparative results and a connection with the split Levinson algorithm are established. Illustrative numerical examples are approximate integrals of the form

$$\int_{-1}^1 \frac{f(x)}{(x+\lambda)^r} \omega(x) dx, \quad r = 1, 2, 3, \dots$$

with  $f(x)$  a continuous function on  $[-1, 1]$ ,  $\omega(x) \geq 0$  a weight function on this interval and  $\lambda \in \mathbb{R}$  such that  $|\lambda| > 1$  is required. Here the classical Gaussian quadrature is an extremely slow procedure.

**Key words.** orthogonal Laurent polynomials, L-Gaussian quadrature, Szegő quadrature, three-term recurrence relations, split Levinson algorithm, numerical quadrature.

**AMS subject classifications.** 41A55, 33C45, 65D30.

**1. Orthogonal Laurent Polynomials.** The important role played by the theory of orthogonal polynomials in the construction of an  $n$ -point quadrature rule

$$(1.1) \quad I_n(f) = \sum_{j=1}^n A_j f(x_j)$$

to approximate integrals of the form

$$(1.2) \quad I_\mu(f) = \int_a^b f(x) d\mu(x), \quad -\infty \leq a < b \leq +\infty$$

( $\mu$  being a positive measure) so that  $I_n(f)$  exactly integrates polynomials of degree as large as possible, is well known. Indeed, let  $\varphi_n(x)$  be the  $n$ -th orthonormal polynomial with respect to the measure  $\mu$  and let  $x_1, \dots, x_n$  be its  $n$  distinct zeros. Then positive weights  $A_1, \dots, A_n$  can be uniquely determined so that the corresponding quadrature formula  $I_n(f)$  satisfies  $I_n(P) = I_\mu(P)$  for all  $P \in \Pi_{2n-1}$  ( $\Pi_k$ : space of polynomials of degree  $k$  at most,  $\Pi$ : space of all polynomials). This is the well known  $n$ -point Gauss-Christoffel (or Gaussian) formula for the measure  $\mu$  on  $(a, b)$  (for a comprehensive survey on this topic see [11]). Furthermore, these formulas are “optimal” in the sense that there is not an  $n$ -point rule (1.1) which is in  $\Pi_{2n}$ . It holds

$$(1.3) \quad A_j = \frac{1}{\sum_{k=0}^{n-1} \varphi_k^2(x_j)}, \quad j = 1, \dots, n.$$

The numerical power and effectiveness of such rules when dealing with smooth integrands  $f(x)$  has been convincingly demonstrated. However, convergence can become extremely slow if the integral has singularities near the integration interval  $(a, b)$ . In order to overcome this drawback, quadrature formulas exactly integrating rational functions with prescribed poles outside  $(a, b)$  have been considered as an alternative in the last years (see e.g. [12], [23] or [3]). Thus, in this paper we shall be concerned with quadrature formulas (1.1) exactly integrating rational functions with all their poles at the origin and the infinity (Laurent polynomials) in order to estimate an integral like  $\int_S f(z) d\mu(z)$ ,  $\mu$  being a

\*Received July 30, 2002. Accepted for publication May 10, 2003. Communicated by J. Arvesú.

†Department of Mathematical Analysis, La Laguna University, 38271 La Laguna. Tenerife. Spain.

positive measure supported on  $S \subset \mathbb{C}$  with either  $S = (a, b)$ ,  $0 \leq a < b \leq +\infty$  or  $S = \mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$  (the unit circle). Here the theory of orthogonal Laurent polynomials will represent the basic ingredient in the construction of the corresponding quadrature rules.

Thus, in general, let  $\mu$  be a positive Borel measure in the complex plane and consider the Hilbert space  $L_2(\mu)$  of measurable functions  $\phi(z)$  for which  $\int |\phi(z)|^2 d\mu(z) < +\infty$ . As usual, in  $L_2(\mu)$  we can define the inner product

$$(1.4) \quad \langle \phi, \psi \rangle = \langle \phi, \psi \rangle_\mu = \int \phi(z) \overline{\psi(z)} d\mu(z), \quad \phi, \psi \in L_2(\mu).$$

Suppose that  $\{\phi_k(z)\}_{k=0}^n$  is a system of linearly independent functions in  $L_2(\mu)$ . By applying the Gram-Schmidt orthogonalization process to  $\{\phi_k(z)\}_{k=0}^n$  a new system of linearly independent functions  $\{\varphi_k(z)\}_{k=0}^n$  can be obtained such that  $\varphi_n(z)$  is a linear combination of the  $n+1$  functions  $\{\phi_k(z)\}_{k=0}^n$  and

$$\langle \varphi_n(z), \varphi_m(z) \rangle = \int \varphi_n(z) \overline{\varphi_m(z)} dz = k_n \delta_{n,m}, \quad k_n > 0.$$

When the process is repeated for each natural  $n$ , an essentially unique system  $\{\varphi_n(z)\}_{n=0}^\infty$  of orthogonal functions with respect to the measure  $\mu$  is obtained. If moreover

$$\|\varphi_n(z)\|^2 = \int |\varphi_n(z)|^2 d\mu(z) = 1,$$

then  $\{\varphi_n(z)\}_{n=0}^\infty$  is called an orthonormal system. For our purposes we will concentrate on the linearly independent system of monomials both with positive and negative exponents, say  $\dots, z^{-2}, z^{-1}, 1, z, z^2, \dots$  yielding the so-called *Laurent polynomials*. Indeed, for  $p$  and  $q$  integers such that  $p \leq q$ ,  $\Delta_{p,q}$  will denote the space of Laurent polynomials of the form  $L(z) = \sum_{j=p}^q \alpha_j z^j$  with  $\alpha_j \in \mathbb{C}$  and  $\Delta$  the space of all Laurent polynomials. Observe that  $\Delta_{0,k} = \Pi_k$ .

Now, in order to generate a sequence of nested subspaces of Laurent polynomials similar to the sequence of subspaces  $\{\Pi_k\}_{k \geq 0}$ , we will start from two nondecreasing sequences of nonnegative integers  $\{p(n)\}_{n=0}^\infty$  and  $\{q(n)\}_{n=0}^\infty$  such that  $p(n) + q(n) = n$  for all  $n = 0, 1, 2, \dots$  and set

$$\mathcal{L}_n = \Delta_{-p(n), q(n)} = \text{Span} \{z^j; -p(n) \leq j \leq q(n)\}.$$

Observe that  $\mathcal{L}_0 = \text{Span}\{1\}$ ,  $\dim(\mathcal{L}_n) = n+1$  and that  $\mathcal{L}_n \subset \mathcal{L}_{n+1}$ . Furthermore if  $p(n) = 0$  then  $q(n) = n$  and  $\mathcal{L}_n = \Delta_{0,n} = \Pi_n$ . In the sequel, in order to guarantee that  $\bigcup_{n=0}^\infty \mathcal{L}_n = \Delta$ , we will assume that  $\lim_{n \rightarrow \infty} p(n) = \lim_{n \rightarrow \infty} q(n) = \infty$  and further restrict ourselves to the particular case

$$(1.5) \quad p(n) = E\left[\frac{n+1}{2}\right], \quad q(n) = n - p(n) = E\left[\frac{n}{2}\right], \quad n \geq 0,$$

where, as usual,  $E[x]$  denotes the integer part of  $x$ . In other words, the sequence  $\{p(n)\}$  has induced the following "ordering" in  $\Delta$ :

$$\Delta_{0,0}, \Delta_{-1,0}, \Delta_{-1,1}, \Delta_{-2,1}, \dots$$

Thus, according to this ordering we can construct a unique sequence (up to a sign)  $\{\varphi_n(z)\}_{n=0}^\infty$  of Laurent polynomials satisfying:

1.  $\varphi_n \in \mathcal{L}_n \setminus \mathcal{L}_{n-1}$ ,  $n = 1, 2, \dots$
2.  $\langle \varphi_n(z), R \rangle_\mu = 0$  for all  $R \in \mathcal{L}_{n-1}$  (i.e.,  $\varphi_n \perp \mathcal{L}_{n-1}$ ).
3.  $\|\varphi_n(z)\|_\mu^2 = \langle \varphi_n(z), \varphi_n(z) \rangle_\mu = 1$ .

In this case,  $\{\varphi_n(z)\}_{n=0}^\infty$  will be called an orthonormal Laurent polynomial sequence corresponding to the measure  $\mu$  and the ordering induced by  $p(n) = E \left[ \frac{n+1}{2} \right]$  which has been the most extensively treated case in the literature. For other choices of  $p(n)$ , see, e.g., [27], [4] or [30]. As already said, in the rest of the paper we will concentrate on measures  $\mu$  supported not on the whole complex plane  $\mathbb{C}$  but either on intervals  $(a, b)$  of the positive real half-line or on the unit circle  $\mathbb{T}$ . In the two following sections the differences and similarities for sequences of orthogonal Laurent polynomials concerning both situations will be displayed. In this respect and in order to fix a uniquely determined sequence of orthogonal Laurent polynomials, we will use the concept of leading coefficient associated with a Laurent polynomial (see [7]). Thus, for a given Laurent polynomial  $R$  there exists a unique natural number  $n$  such that  $R \in \mathcal{L}_n$  and  $R \notin \mathcal{L}_{n-1}$ ;  $n$  is called the “L-degree” of  $R$  for the ordering established above. Observe that if  $P(z)$  is a polynomial of degree  $n$ , then its L-degree is just  $2n$ . Indeed,  $P(z) \in \Delta_{-n,n} = \mathcal{L}_{2n}$  and  $P(z) \notin \mathcal{L}_{2n-1}$ . Setting  $R(z) = \sum_{j=-p(n)}^{q(n)} \alpha_j z^j$ , one defines the leading coefficient of  $R$  as  $\alpha_{q(n)}$  if  $n$  is even or  $\alpha_{-p(n)}$  if  $n$  is odd. Clearly, for an orthogonal sequence  $\{\phi_n(z)\}_{n=0}^\infty$  of Laurent polynomials,  $\phi_n(z)$  has a non-zero leading coefficient for each  $n$ , which will be sometimes assumed positive. When equal to one,  $\phi_n(z)$  is said monic. Clearly, an orthonormal sequence  $\{\varphi_n(z)\}$  of Laurent polynomials with positive leading coefficient for each  $n$  is uniquely determined.

The paper is organized as follows: In Section 2,  $n$ -point quadrature formulas for a measure  $\mu$  on  $S = (a, b)$ ,  $0 \leq a < b \leq \infty$ , exactly integrating Laurent polynomials belonging to certain subspaces of dimension  $2n$  are characterized. A three-term recurrence relation for the sequence of orthogonal Laurent polynomials and a Christoffel-Darboux formula are also mentioned. The same problem is studied in Section 3 for the unit circle  $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$  and unlike the situation in Section 2, it is proved that there can not exist an  $n$ -point quadrature formula with nodes on  $\mathbb{T}$  that is exact in an appropriate subspace of  $\Delta$  of dimension  $2n$ . The corresponding formulas with highest degree of exactness in this situation ( $2n - 1$ ) are characterized and a three-term recurrence relation for the sequence of monic orthogonal Laurent polynomials on the unit circle is proved. In Section 4 we establish that the monic orthogonal Laurent polynomials are closely related to the first and second singular predictor polynomials computed recursively for the split Levinson algorithm. Finally, in Section 5 illustrative numerical examples are given.

**2. The Real Half-Line: L-Gaussian Quadratures.** In this section we will assume that the measure  $\mu$  is supported on the interval  $(a, b)$  where  $0 \leq a < b \leq +\infty$ . Thus, we are interested in approximating

$$(2.1) \quad I_\mu(f) = \int_a^b f(x) d\mu(x)$$

by means of an  $n$ -point quadrature rule

$$(2.2) \quad I_n(f) = \sum_{j=1}^n A_j f(x_j)$$

where the nodes (which are assumed distinct and in  $(a, b)$ ) and weights are to be determined by imposing

$$(2.3) \quad I_\mu(R) = I_n(R)$$

for any Laurent polynomial  $R$  with L-degree as large as possible, i.e.,  $R \in \mathcal{L}_N$ , where  $N = N(n)$  and as large as possible. For this purpose, proceed in a way similar to the polynomial case. Thus, since  $\mathcal{L}_{n-1}$  is a Tchebyshev space of dimension  $n$  on any subinterval of  $(a, b)$ , given  $n$  distinct nodes  $x_1, x_2, \dots, x_n$  on  $(a, b)$ , the weights  $A_1, \dots, A_n$  can be determined so that

$$(2.4) \quad I_\mu(R) = I_n(R) = \sum_{j=1}^n A_j f(x_j), \quad \forall R \in \mathcal{L}_{n-1}.$$

Furthermore, it also holds that if  $L_{n-1}(f, \cdot)$  represents the unique Laurent polynomial in  $\mathcal{L}_{n-1}$  interpolating  $f$  at the nodes  $\{x_j\}_{j=1}^n$ , then

$$(2.5) \quad I_n(f) = I_\mu(L_{n-1}(f, \cdot)).$$

For this reason  $I_n(f)$  given by (2.4) sometimes will be called of interpolatory type.

Now, in order to increase the ‘‘L-degree’’ of exactness we have the following result.

**THEOREM 2.1.** *Let  $I_n(f) = \sum_{j=1}^n A_j f(x_j)$  be an  $n$ -point quadrature formula such that  $x_i \neq 0$  for  $i = 1, \dots, n$ . Then,  $I_n(f) = I_\mu(f)$  for all  $f \in \mathcal{L}_{n+r}$  with  $r \geq 0$  if and only if*

1.  $I_n(f)$  is of interpolatory type
- 2.

$$(2.6) \quad \langle R_n(x), h(x) \rangle = 0, \quad \forall h(x) \in \mathcal{L}_r \quad (\text{i.e., } R_n(x) \perp \mathcal{L}_r),$$

$$\text{where } R_n(x) = \frac{Q_n(x)}{x^{p(n)}} = \frac{\prod_{j=1}^n (x - x_j)}{x^{p(n)}} \in \mathcal{L}_n.$$

*Proof.* Similar to the proof of Theorem 3.1 (in the next section). See also [20] for the polynomial situation with  $r = n - 1$ . □

It should be taken into account that since we are dealing with real-valued functions, we do not need complex conjugates in the inner product. On the other hand and concerning the integer  $r$ , one actually has  $0 \leq r \leq n - 1$ , since it is very easy to check that there does not exist an  $n$ -point quadrature formula which is exact in  $\mathcal{L}_N$  with  $N \geq 2n$ . Thus, we are interested in constructing quadrature formulas, like (2.2), exact in  $\mathcal{L}_{n+r}$  with  $0 \leq r \leq n - 1$ . From (2.6) it follows that

$$(2.7) \quad R_n(z) = \sum_{j=r+1}^n \alpha_j \varphi_j(z),$$

where  $\{\varphi_n(z)\}_{n=0}^\infty$  is the corresponding sequence of orthonormal Laurent polynomials for the measure  $\mu$  on  $(a, b)$ . Now, the following question immediately arises: Is it possible to choose the parameters  $\alpha_j$  in (2.7) so that  $R_n(z)$  has exactly  $n$  distinct zeros in  $(a, b)$ ? A positive response can be given when the largest domain of exactness is required, i.e.  $r = n - 1$ .

**THEOREM 2.2.** *Let  $\varphi_n(x)$  denote the  $n$ -th orthonormal Laurent polynomial with respect to the measure  $\mu$  on  $(a, b)$ . Then,*

1.  $\varphi_n(x)$  has exactly  $n$  distinct zeros on  $(a, b)$ .
2. The zeros of  $\varphi_n(x)$  and  $\varphi_{n+1}(x)$  interlace.
3. Let  $x_1, \dots, x_n$  be the zeros of  $\varphi_n(x)$ . Then there exist positive weights  $A_1, \dots, A_n$ , such that

$$(2.8) \quad I_n(f) = \sum_{j=1}^n A_j f(x_j) = I_\mu(f) = \int_a^b f(x) d\mu(x), \quad \forall f \in \mathcal{L}_{2n-1}.$$

*Proof.* See [7], [19] or [16]. □

REMARK 2.3. The quadrature formulas given by (2.8) were earlier introduced in [19] (see also [18]). Through this last paper a new area of mathematics was opened leading to a strong moment problem and related topics (continued fractions, two-point Padé approximants, . . .) and where orthogonal Laurent polynomials made their appearance. These quadrature rules which we shall refer to as L-Gaussian formulas, have been considered by different authors in the last years (see [7], [4] or [5]). For alternative approaches see [26] and [22]. As for numerical experiments see [26], [2] and [15].

In the rest of the section we will survey the most relevant features of the L-Gaussian formulas in connection with the orthogonal Laurent polynomials. The closest parallelism with classical Gaussian formulas and orthogonal polynomials will be specially emphasized. We start with a three-term recurrence relation (see [26]):

THEOREM 2.4. Let  $\{R_n(z)\}_{n=0}^\infty$  be the sequence of orthogonal Laurent polynomials normalized as follows. Set  $R_n(z) = \frac{B_n(z)}{z^{p(n)}}$ , where  $B_n(z)$  is a monic polynomial of degree  $n$ . Then

$$(2.9) \quad R_{n+1}(z) = (z - \beta_{n+1})z^{d(n)}R_n(z) - \alpha_{n+1}R_{n-1}(z), \quad n \geq 0,$$

with  $R_{-1}(z) \equiv 0$ ,  $R_0(z) \equiv 1$ , where

$$d(n) = \begin{cases} -1 & \text{if } n \text{ is even} \\ 0 & \text{if } n \text{ is odd} \end{cases}$$

and

$$\alpha_1 = \mu_0, \quad \beta_1 = \frac{\mu_0}{\mu_{-1}}, \quad \alpha_{n+1} = \frac{\rho_n}{\rho_{n-1}}, \quad \beta_{n+1} = -\alpha_{n+1} \frac{\sigma_{n-1}}{\sigma_n}, \quad n \geq 1.$$

Here,

$$\mu_k = \int_a^b x^k d\mu(x), \quad \rho_n = \langle R_n, t^{p(n)} \rangle, \quad \sigma_n = \langle R_n, t^{-(q(n)+1)} \rangle.$$

(Recall  $p(n) = E \left[ \frac{n+1}{2} \right]$ ,  $q(n) = E \left[ \frac{n}{2} \right]$ ). Furthermore, both  $\alpha_n$  and  $\beta_n$  are positive for  $n \geq 1$ . □

REMARK 2.5. In [7] a Favard-type theorem for the sequence  $\{R_n(z)\}_{n=0}^\infty$  is proved.

For the weights  $A_1, \dots, A_n$  in the  $n$ -point L-Gaussian formula, similar results to the polynomial case are also true. Since  $\varphi_n \in \mathcal{L}_n = \Delta_{-p(n), q(n)}$  and has  $n$  distinct zeros on  $(0, \infty)$  we can write  $\varphi_n(x) = v_n x^{-p(n)} + \dots + u_n x^{q(n)}$  ( $u_n v_n \neq 0$ ). We will also require the normalization conditions:  $u_n > 0$  for all  $n = 0, 1, 2, \dots$ . First, we need the following Christoffel-Darboux formula (see [24]):

THEOREM 2.6 (Christoffel-Darboux). Let  $\{\varphi_n(x)\}_{n=0}^\infty$  be the orthonormal sequence of Laurent polynomials. Then, setting  $\lambda(n) = (-1)^n$ , one has

(2.10)

$$\sum_{i=0}^n \varphi_i(x)\varphi_i(y) = \frac{\lambda(n)v_n}{u_{n+1}} \left[ \frac{(xy)^{1/2} \left( \left( \frac{x}{y} \right)^{\lambda(n)} \varphi_{n+1}(x)\varphi_n(y) - \left( \frac{y}{x} \right)^{\lambda(n)} \varphi_n(x)\varphi_{n+1}(y) \right)}{x - y} \right]$$

□

Let  $l_j \in \mathcal{L}_{n-1}$  be such that  $l_j(x_k) = \delta_{j,k}$ , where  $\{x_1, \dots, x_n\}$  as usual are the zeros of the  $n$ -th orthonormal Laurent polynomial  $\varphi_n(x)$ . Clearly  $l_j^2 \in \mathcal{L}_{2n-1}$ , hence  $A_j = I_\mu(l_j^2) > 0$ ,  $j = 1, \dots, n$ . On the other hand, from (2.10) with  $x = x_j$  and letting  $y$  tend to  $x_j$ , the following theorem can be proved (see [6] for details):

**THEOREM 2.7.** *Let  $\varphi_n(x)$  be the  $n$ -th orthonormal Laurent polynomial for the measure  $\mu$ . Then, the weights  $\{A_j\}_{j=1}^n$  are given by*

$$A_j = \frac{1}{\sum_{i=0}^{n-1} \varphi_i^2(x_j)}, \quad j = 1, \dots, n.$$

**REMARK 2.8.** *As a conclusion we can say that the  $n$ -point  $L$ -Gaussian quadrature rule (1.1) is completely characterized in terms of the orthonormal Laurent polynomials  $\{\varphi_k(z)\}_{k=0}^n$  which can be recursively computed by relation (2.9).*

**3. The Unit Circle: Szegő Quadratures.** In this section we shall be concerned with positive measures supported on the unit circle  $\mathbb{T}$  which in the sequel will be denoted by  $\sigma$  (in order to avoid a possible confusion with the measure  $\mu$  on  $(0, \infty)$  considered in Section 2). Since we will deal now with complex-valued functions, complex conjugation will be again required in the inner product induced by  $\sigma$ , i.e.

$$(3.1) \quad \langle f, g \rangle_\sigma = \int_{-\pi}^{\pi} f(e^{i\theta}) \overline{g(e^{i\theta})} d\sigma(\theta), \quad f, g \in L_2^\sigma(\mathbb{T})$$

We are interested in approximating the integral

$$(3.2) \quad I_\sigma(f) = \int_{\mathbb{T}} f(z) d\sigma(z) = \int_{-\pi}^{\pi} f(e^{i\theta}) d\sigma(\theta)$$

by an  $n$ -point quadrature rule with nodes on  $\mathbb{T}$ ,

$$(3.3) \quad I_n(f) = \sum_{j=1}^n \lambda_j f(z_j), \quad z_j \neq z_k \quad (j \neq k), \quad z_j \in \mathbb{T}, \quad \forall j = 1, \dots, n$$

Thus, starting from  $n$  distinct nodes  $z_1, \dots, z_n$  on  $\mathbb{T}$ , and since  $\mathcal{L}_{n-1}$  is also a Tchebyshev space of dimension  $n$  on  $\mathbb{T}$ , weights  $\lambda_1, \dots, \lambda_n$  can be determined so that

$$(3.4) \quad I_\sigma(f) = I_n(f), \quad \forall f \in \mathcal{L}_{n-1}.$$

Here it also holds that  $I_n(f)$ , characterized by (3.4), can be represented as

$$(3.5) \quad I_n(f) = I_\sigma(R_{n-1}(f, \cdot)),$$

where  $R_{n-1}(f, \cdot)$  is the Laurent polynomial in  $\mathcal{L}_{n-1}$  which interpolates  $f$  at the nodes  $\{z_j\}_{j=1}^n$ . However, the first difference with the real half-line appears when an appropriate selection of the nodes  $\{z_j\}_{j=1}^n$  is required in order to increase the dimension of the subspace of Laurent polynomials where exactness of the quadrature formula takes place. Because of the inner product (3.1), we need the following subspaces for  $r \geq 0$ ,

$$\mathcal{L}_{r*} = \{f \in \Delta, f_* \in \mathcal{L}_r\} = \Delta_{-q(r), p(r)},$$

where for  $f \in \Delta$ :  $f_*(z) = \overline{f\left(\frac{1}{\bar{z}}\right)}$  (“substar-conjugation”). Under these conditions, orthogonality immediately arises, as follows (compare with Theorem 2.1).

**THEOREM 3.1.** *Let  $I_n(f) = \sum_{j=1}^n \lambda_j f(z_j)$  be an  $n$ -point quadrature formula such that  $z_j \neq 0$  for  $j = 1, \dots, n$ . Then  $I_n(f)$  is exact in  $\mathcal{L}_n \mathcal{L}_{r*}$ ,  $r \geq 0$ , if and only if*

1.  $I_n(f)$  is exact in  $\mathcal{L}_{n-1}$ .
- 2.

$$(3.6) \quad \langle R_n(z), g(z) \rangle_\sigma = 0, \quad \forall g \in \mathcal{L}_r \quad (\text{i.e., } R_n \perp \mathcal{L}_r),$$

$$\text{where } R_n(z) = \frac{Q_n(z)}{z^{p(n)}} = \frac{\prod_{j=1}^n (z - z_j)}{z^{p(n)}} \in \mathcal{L}_n.$$

*Proof.*- “ $\implies$ ” 1.- Trivial. 2.- Recall that  $\mathcal{L}_n \mathcal{L}_{r^*} = \Delta_{-(p(n)+q(r)), (q(n)+p(r))}$  and  $\mathcal{L}_r = \Delta_{-p(r), q(r)}$ . Thus,  $\langle R_n(z), g(z) \rangle_\sigma = 0$  for all  $g(z) \in \mathcal{L}_r$  is equivalent to  $\langle R_n(z), z^j \rangle_\sigma = 0$  for all  $-p(r) \leq j \leq q(r)$ . Now,

$$\langle R_n(z), z^j \rangle_\sigma = \int_{-\pi}^{\pi} R_n(z) z^{-j} \sigma(\theta) d\theta = \int_{-\pi}^{\pi} \frac{Q_n(z)}{z^{p(n)+j}} \sigma(\theta) d\theta.$$

For  $j = q(r)$  and since  $Q_n(z)$  has exact degree  $n$ , we see that

$$\begin{aligned} \frac{Q_n(z)}{z^{p(n)+q(r)}} &\in \Delta_{-(p(n)+q(r)), (n-p(n)-q(r))} = \Delta_{-(p(n)+q(r)), (q(n)-q(r))} \subset \\ &\subset \Delta_{-(p(n)+q(r)), (q(n)+q(r))} = \mathcal{L}_n \mathcal{L}_{r^*}. \end{aligned}$$

On the other hand, for  $j = -p(r)$  we have

$$\begin{aligned} \frac{Q_n(z)}{z^{p(n)-p(r)}} &\in \Delta_{-(p(n)-p(r)), (n-p(n)-p(r))} = \Delta_{-(p(n)-p(r)), (q(n)+p(r))} \subset \\ &\subset \Delta_{-(p(n)+q(r)), (q(n)+p(r))} = \mathcal{L}_n \mathcal{L}_{r^*}. \end{aligned}$$

Thus, for  $-p(r) \leq j \leq q(r)$ ,  $R_n(z) z^{-j} \in \mathcal{L}_n \mathcal{L}_{r^*}$ . Therefore one can write

$$\langle R_n(z), z^j \rangle_\sigma = I_\sigma (R_n(z) z^{-j}) = I_n (R_n(z) z^{-j}) = 0$$

since  $R_n(z_j) = 0$ .

“ $\Leftarrow$ ” Let  $z_1, \dots, z_n$  be the zeros of  $R_n(z) \in \mathcal{L}_n$  so that (3.6) holds. Then a quadrature formula based upon these nodes, such as  $I_n(f) = \sum_{j=1}^n \lambda_j f(z_j)$ , can be determined so that it is exact in  $\mathcal{L}_{n-1}$ . Setting  $L^{(j)}(z) \in \mathcal{L}_{n-1}$  such that  $L^{(j)}(z_k) = \delta_{jk}$ , then  $\lambda_j = I_\sigma(L^{(j)})$  for all  $j = 1, \dots, n$ .

Take  $L(z) \in \mathcal{L}_n \mathcal{L}_{r^*}$  and define  $A(z) = L(z) - \sum_{j=1}^n L(z_j) L^{(j)}(z)$ . Thus,  $A \in \mathcal{L}_n \mathcal{L}_{r^*}$ ,  $A(z_j) = 0$  for all  $j = 1, \dots, n$ , and we can write

$$A(z) = \frac{Q_n(z) S(z)}{z^{p(n)+q(r)}}, \quad S(z) \in \Pi_r.$$

Observe that  $A(z) = R_n(z) H(z)$  with  $H(z) = \frac{S(z)}{z^{q(r)}}$ . Hence, setting  $T(z) = H_*(z) \in \mathcal{L}_r$ , we have

$$I_\sigma(A(z)) = I_\sigma(R_n(z) T_*(z)) = \langle R_n(z), T(z) \rangle_\sigma = 0,$$

which yields

$$\begin{aligned} I_\sigma(L(z)) &= I_\sigma \left( A(z) + \sum_{j=1}^n L(z_j) L^{(j)}(z) \right) = I_\sigma(A(z)) + \\ &+ I_\sigma \left( \sum_{j=1}^n L(z_j) L^{(j)}(z) \right) = \sum_{j=1}^n I_\sigma(L^{(j)}(z)) L(z_j) = I_n(L). \end{aligned}$$

□

Consider now the family of monic Szegő polynomials  $\{\rho_n(z)\}_{n=0}^\infty$  orthogonal with respect to the inner product (3.1), i.e., for  $n = 0, 1, \dots$  then  $\rho_n(z)$  satisfies

$$(3.7) \quad \langle \rho_n(z), z^m \rangle_\sigma = 0 \quad (0 \leq m \leq n-1)$$

and

$$(3.8) \quad \langle \rho_n(z), \rho_n(z) \rangle_\sigma = \langle \rho_n(z), z^n \rangle_\sigma = \Delta_n / \Delta_{n-1},$$

where  $\Delta_n$  is the  $n$ -th Toeplitz determinant of the trigonometric moments of  $\sigma$ . Set  $\rho_n^*(z) = z^n \rho_n(z)$  (reciprocal polynomial). Then, from the above orthogonality conditions both for  $\{\rho_n(z)\}_{n=0}^\infty$  and  $\{\phi_n(z)\}_{n=0}^\infty$  (recall that  $\phi_n(z)$  denotes the  $n$ -th orthogonal Laurent polynomial with respect to the measure  $\sigma$ ) the following relation between  $\phi_n(z)$  and  $\rho_n(z)$  is deduced:

$$(3.9) \quad \phi_{2n}(z) = \frac{1}{z^n} \rho_{2n}(z), \quad \phi_{2n+1}(z) = \frac{1}{z^{n+1}} \rho_{2n+1}^*(z).$$

On the other hand, it is well known (see [1]) that the zeros of the polynomial  $\rho_n(z)$  are located in  $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$  and consequently, the zeros of  $\rho_n^*(z)$  in  $\mathbb{E} = \{z \in \mathbb{C} : |z| > 1\}$ . From this fact and (3.9) we obtain the following result

LEMMA 3.2. *The zeros of  $\phi_{2n}(z)$  and  $\phi_{2n+1}(z)$  are located in  $\mathbb{D}$  and  $\mathbb{E}$  respectively.*  $\square$

REMARK 3.3. *Actually  $r$  in Theorem 3.1 should be taken such that  $0 \leq r \leq n-1$ . Indeed, assume  $r \geq n$ , i.e.,  $r = n+k$  with  $k \geq 0$ . Then,  $\mathcal{L}_n \mathcal{L}_{r*} = \Delta_{-(n+q(k)), (n+p(k))}$  since for  $p(n) = E\left[\frac{n+1}{2}\right]$  and  $q(n) = n - p(n) = E\left[\frac{n}{2}\right]$  it holds that  $p(r+s) = p(r) + p(s)$  and  $q(r+s) = q(r) + q(s)$ . Thus, if  $Q_n(z) = \prod_{j=1}^n (z - z_j)$ , then*

$$|Q_n(z)|^2 = Q_n(z) \overline{Q_n(z)} \in \Delta_{-n,n} \subset \mathcal{L}_n \mathcal{L}_{r*} \quad (z = e^{i\theta}).$$

Hence,  $0 < \int_{-\pi}^{\pi} |Q_n(z)|^2 d\sigma(\theta)$  and  $I_n(|Q_n(z)|^2) = 0$ .

Let us next see what happens with the largest reachable domain of exactness, i.e., if  $r = n-1$ . Unlike the situation on the half-line we have the following negative result.

THEOREM 3.4. *There can not exist an  $n$ -point quadrature formula with nodes on  $\mathbb{T}$  to be exact in  $\mathcal{L}_n \mathcal{L}_{(n-1)*}$ . Proof.- Suppose that there exists such a formula:*

$$I_n(f) = \sum_{j=1}^n \lambda_j f(z_j) = I_\sigma(f)$$

where  $|z_j| = 1$  for  $j = 1, \dots, n$ . Define  $R_n(z) = \frac{(z-z_1)\cdots(z-z_n)}{z^{p(n)}} \in \mathcal{L}_n$ . Then by Theorem 3.1,  $R_n \perp \mathcal{L}_{n-1}$ .

Thus, if  $\{\phi_k(z)\}_{k=0}^\infty$  is a sequence of orthogonal Laurent polynomials, then  $R_n(z) = \lambda_n \phi_n(z)$  ( $\lambda_n \neq 0$ ). So, the nodes  $z_1, \dots, z_n$  are the zeros of  $\phi_n(z)$ , which can not be located on  $\mathbb{T}$  by Lemma 3.2 and a contradiction arises.  $\square$

Now, the next step would be to consider the case  $r = n-2$ . More precisely, one might wonder is it possible to construct an  $n$ -point quadrature formula with distinct nodes  $\{z_j\}$  on  $\mathbb{T}$  to be exact in  $\mathcal{L}_n \mathcal{L}_{(n-2)*}$ ? By Theorem 3.1 if  $R_n(z) = \frac{\prod_{j=1}^n (z-z_j)}{z^{p(n)}}$  then  $R_n(z) \perp \mathcal{L}_{n-2}$ . Hence,

$$(3.10) \quad R_n(z) = \alpha \phi_n(z) + \beta \phi_{n-1}(z), \quad \alpha, \beta \in \mathbb{C},$$



where  $\{\phi_k(z)\}_{k=0}^{\infty}$  is a sequence of orthogonal Laurent polynomials. So, from (3.10) the above question can be reformulated as follows: Is it possible to find  $\alpha$  and  $\beta$  in (3.10) so that  $R_n(z)$  has its zeros on  $\mathbb{T}$  ?

**THEOREM 3.5.** *Parameters  $\alpha$  and  $\beta$  in (3.10) can be conveniently chosen so that  $R_n(z) = \alpha\phi_n(z) + \beta\phi_{n-1}(z)$  has exactly  $n$  distinct zeros on  $\mathbb{T}$ .*

*Proof.* Setting  $R_n(z) = \frac{N_n(z)}{z^{p(n)}}$ , with  $N_n(z)$  being a polynomial of degree  $n$  at most depending on  $\alpha$  and  $\beta$ . These parameters should be chosen so that  $N_n(z)$  satisfies the following:

1.  $N_n(z)$  has exact degree  $n$ .
2.  $N_n^*(z) = \lambda_n N_n(z)$  with  $\lambda_n \neq 0$ .
3.  $\langle N_n(z), z^k \rangle_{\sigma} = 0, 1 \leq k \leq n-1, \langle N_n(z), 1 \rangle_{\sigma} \neq 0, \langle N_n(z), z^n \rangle_{\sigma} \neq 0$ .

Now by using Theorem 6.2 in [17] the proof follows. □

Hence, from Theorems 3.1 and 3.5 we obtain the the following result

**THEOREM 3.6.** *Let  $\{z_1, \dots, z_n\}$  be the  $n$  distinct zeros of  $R_n(z) = \alpha\phi_n(z) + \beta\phi_{n-1}(z)$  as given in Theorem 3.5. Then, there exist positive numbers  $\lambda_1, \dots, \lambda_n$  such that*

$$I_n(f) = \sum_{j=1}^n \lambda_j f(z_j) = I_{\sigma}(f) \quad , \quad \forall f \in \mathcal{L}_n \mathcal{L}_{(n-2)*}.$$

*Proof.* It only remains to prove that the weights  $\lambda_1, \dots, \lambda_n$  are positive. Indeed, for  $1 \leq j \leq n$ , let  $L^{(j)}(z) \in \mathcal{L}_{(n-1)}$  be such that  $L_j(z_k) = \delta_{j,k}$  for  $1 \leq j, k \leq n$ . Then, for  $z \in \mathbb{T}$ ,  $|L_j(z)|^2 \in \mathcal{L}_{(n-1)} \mathcal{L}_{(n-1)*} = \mathcal{L}_n \mathcal{L}_{(n-2)*}$  as can be checked easily. Now, the proof immediately follows, since

$$0 < I_{\sigma}(|L_j|^2) = \sum_{k=1}^n \lambda_k |L_j(z_k)|^2 = \lambda_j \quad , \quad j = 1, \dots, n.$$

□

So, from Theorems 3.1 and 3.6 quadrature formulas with nodes on  $\mathbb{T}$  which are exact in  $\mathcal{L}_n \mathcal{L}_{(n-2)*}$  can be essentially characterized by means of the Laurent polynomials of the form

$$(3.11) \quad R_n(z) = \alpha\phi_n(z) + \beta\phi_{n-1}(z),$$

which by analogy with the polynomial situation (see e.g. [1]) could be called “para-orthogonal Laurent polynomials”. On the other hand, one also has that

$$\mathcal{L}_n \mathcal{L}_{(n-2)*} = \mathcal{L}_{(n-1)} \mathcal{L}_{(n-1)*} = \Delta_{-(n-1), (n-1)}.$$

Quadrature formulas with nodes on  $\mathbb{T}$  to be exact in  $\Delta_{-(n-1), (n-1)}$  were earlier introduced by Jones, et al. ([17]) in connection with the solution of the trigonometric moment problem (see also [29] and [14]). Such quadratures were named “Szegő formulas” and all studied in a series of recent papers: [13], [25], [8] [9].

As a consequence, making  $R_n(z) = \alpha\phi_n(z) + \beta\phi_{n-1}(z) = \frac{N_n(z)}{z^{p(n)}}$ , by Theorem 6.1 in [17] one can write

$$N_n(z) = \lambda_n [\rho_n(z) + \tau_n \rho_n^*(z)] \quad , \quad \lambda_n \neq 0 \quad , \quad |\tau_n| = 1.$$

In [17] polynomials of the form  $\rho_n(z) + \tau \rho_n^*(z)$  are called “para-orthogonal”. In the next section we will concentrate on the particular cases  $\tau_n = \pm 1$ .

In the rest of this section we will continue to emphasize the analogies and differences between the real half-line and the unit circle. The following theorem yields an alternative expression for the weights  $\lambda_j$  (compare with Theorem 2.7).

**THEOREM 3.7.** *Let  $\{\tilde{\phi}_k\}_{k=0}^{\infty}$  be the sequence of orthonormal Laurent polynomials and set  $I_n(f) = \sum_{j=1}^n \lambda_j f(z_j) = I_{\sigma}(f)$  for all  $f \in \mathcal{L}_n \mathcal{L}_{(n-2)*}$ . Then*

$$\lambda_j = \frac{1}{\sum_{k=0}^{n-1} |\tilde{\phi}_k(z_j)|^2}.$$

*Proof.* Direct consequence of Proposition 3.4 in [13] and (3.9). □

**REMARK 3.8.** *Observe that unlike the real half-line situation now the nodes  $\{z_j\}_{j=1}^n$  in the quadrature formula are not the zeros of  $\tilde{\phi}_n(z)$*

**THEOREM 3.9 (Three-term recurrence relation).** *Let  $\{\delta_n\}_{n=1}^{\infty}$  be the sequence of Schur parameters (or reflection coefficients) for the measure  $\sigma$ , i.e.,  $\delta_n = \rho_n(0)$ , where  $\rho_n(z)$  is the  $n$ -th monic Szegő polynomial with respect to the measure  $\sigma$ , and let  $\{\phi_n(z)\}_{n=0}^{\infty}$  be the sequence of monic orthogonal Laurent polynomials. Then, it holds that*

$$(3.12) \quad \phi_n(z) = \left( A_n + \overline{A_{n-1}} z^{(-1)^n} \right) \phi_{n-1}(z) + (1 - |\delta_{n-1}|^2) z^{(-1)^n} \phi_{n-2}(z), \quad n \geq 2,$$

with

$$\phi_0(z) \equiv 1, \quad \phi_1(z) = \overline{\delta_1} + \frac{1}{z}, \quad A_n = \begin{cases} \delta_n, & \text{if } n \text{ is even,} \\ \overline{\delta_n}, & \text{if } n \text{ is odd.} \end{cases}$$

*Proof.*- From the recurrence relations satisfied by the Szegő polynomials (see [28]):

$$(3.13) \quad \left. \begin{aligned} \rho_0(z) &= \rho_0^*(z) \equiv 1 \\ \rho_n(z) &= z\rho_{n-1}(z) + \delta_n \rho_{n-1}^*(z) \\ \rho_n^*(z) &= \overline{\delta_n} z \rho_{n-1}(z) + \rho_{n-1}^*(z) \end{aligned} \right\} n = 1, 2, \dots$$

we deduce the initial conditions  $\phi_0(z) = \rho_0(z) \equiv 1$ ,  $\phi_1(z) = \frac{1}{z} \rho_1^*(z) = \frac{1}{z} (1 + \overline{\delta_1} z) = \overline{\delta_1} + \frac{1}{z}$  (since  $\rho_1(z) = \rho_1(0) + z = \delta_1 + z$ ) and the relations

$$(3.14) \quad \begin{aligned} \phi_{2n}(z) &= \frac{1}{z^n} \rho_{2n}(z) = \frac{1}{z^n} [z\rho_{2n-1}(z) + \delta_{2n} \rho_{2n-1}^*(z)] = \frac{1}{z^{n-1}} \rho_{2n-1}(z) + \\ &+ \delta_{2n} \frac{1}{z^n} \rho_{2n-1}^*(z) = \frac{1}{z^{n-1}} [z\rho_{2n-2}(z) + \delta_{2n-1} \rho_{2n-2}^*(z)] + \delta_{2n} \frac{1}{z^n} \rho_{2n-1}^*(z) = \\ &= \frac{1}{z^{n-2}} \rho_{2n-2}(z) + \delta_{2n-1} \frac{1}{z^{n-1}} \rho_{2n-2}^*(z) + \delta_{2n} \frac{1}{z^n} \rho_{2n-1}^*(z) = \frac{1}{z^{n-2}} \rho_{2n-2}(z) + \\ &+ \delta_{2n-1} \frac{1}{z^{n-1}} [\rho_{2n-1}^*(z) - \overline{\delta_{2n-1}} z \rho_{2n-2}(z)] + \delta_{2n} \frac{1}{z^n} \rho_{2n-1}^*(z) = \\ &= \frac{1}{z^{n-2}} \rho_{2n-2}(z) + \delta_{2n-1} \frac{1}{z^{n-1}} \rho_{2n-1}^*(z) - \delta_{2n-1} \overline{\delta_{2n-1}} \frac{1}{z^{n-2}} \rho_{2n-2}(z) + \\ &+ \delta_{2n} \frac{1}{z^n} \rho_{2n-1}^*(z) = (\delta_{2n} + \delta_{2n-1} z) \frac{1}{z^n} \rho_{2n-1}^*(z) + (1 - |\delta_{2n-1}|^2) \frac{1}{z^{n-2}} \rho_{2n-2}(z) \\ &= (\delta_{2n} + \delta_{2n-1} z) \phi_{2n-1}(z) + (1 - |\delta_{2n-1}|^2) z \phi_{2n-2}(z) \end{aligned}$$

and

(3.15)

$$\begin{aligned}
 \phi_{2n+1}(z) &= \frac{1}{z^{n+1}} \rho_{2n+1}^*(z) = \frac{1}{z^{n+1}} \left[ \overline{\delta_{2n+1}} z \rho_{2n}(z) + \rho_{2n}^*(z) \right] = \overline{\delta_{2n+1}} \frac{1}{z^n} \rho_{2n}(z) + \\
 &+ \frac{1}{z^{n+1}} \rho_{2n}^*(z) = \overline{\delta_{2n+1}} \frac{1}{z^n} \rho_{2n}(z) + \frac{1}{z^{n+1}} \left( \overline{\delta_{2n}} z \rho_{2n-1}(z) + \rho_{2n-1}^*(z) \right) = \\
 &= \overline{\delta_{2n+1}} \frac{1}{z^n} \rho_{2n}(z) + \overline{\delta_{2n}} \frac{1}{z^n} \rho_{2n-1}(z) + \frac{1}{z^{n+1}} \rho_{2n-1}^*(z) = \overline{\delta_{2n+1}} \frac{1}{z^n} \rho_{2n}(z) \\
 &+ \overline{\delta_{2n}} \frac{1}{z^{n+1}} \left( \rho_{2n}(z) - \delta_{2n} \rho_{2n-1}^*(z) \right) + \frac{1}{z^{n+1}} \rho_{2n-1}^*(z) = \overline{\delta_{2n+1}} \frac{1}{z^n} \rho_{2n}(z) \\
 &+ \overline{\delta_{2n}} \frac{1}{z^{n+1}} \rho_{2n}(z) - \delta_{2n} \overline{\delta_{2n}} \frac{1}{z^{n+1}} \rho_{2n-1}^*(z) + \frac{1}{z^{n+1}} \rho_{2n-1}^*(z) = \\
 &= \left( \overline{\delta_{2n+1}} + \frac{\overline{\delta_{2n}}}{z} \right) \frac{1}{z^n} \rho_{2n}(z) + (1 - |\delta_{2n}|^2) \frac{1}{z^{n+1}} \rho_{2n-1}^*(z) = \\
 &= \left( \overline{\delta_{2n+1}} + \frac{\overline{\delta_{2n}}}{z} \right) \phi_{2n}(z) + (1 - |\delta_{2n}|^2) \frac{1}{z} \phi_{2n-1}(z).
 \end{aligned}$$

Now (3.12) follows from (3.14) and (3.15).  $\square$

REMARK 3.10. For an alternative proof of Theorem 3.9 based upon certain continued fractions, see [30]. On the other hand, if the trigonometric moments  $\mu_k = \int_{-\pi}^{\pi} e^{-ik\theta} d\sigma(\theta)$  are real, then  $\delta_k \in \mathbb{R}$  and (3.12) can be written as

$$(3.16) \quad \phi_n(z) = \left( \delta_n + \delta_{n-1} z^{(-1)^n} \right) \phi_{n-1}(z) + (1 - |\delta_{n-1}|^2) z^{(-1)^n} \phi_{n-2}(z), \quad n \geq 2.$$

EXAMPLE 3.11. Consider the biparametric family of measures

$$d\sigma(\theta) = \left| \sin\left(\frac{\theta}{2}\right) \right|^{2\alpha+1} \left| \cos\left(\frac{\theta}{2}\right) \right|^{2\beta+1} d\theta, \quad \alpha, \beta > -1.$$

In this case it is known (see [28]) that

$$\delta_{2n} = \frac{\alpha + \beta + 1}{2n + \alpha + \beta + 1} \in \mathbb{R}, \quad \delta_{2n+1} = \frac{\alpha - \beta}{2n + \alpha + \beta + 2} \in \mathbb{R}.$$

When  $\beta = -\frac{1}{2}$ , the measure and the Schur parameters are given by

$$d\sigma(\theta) = \left| \sin\left(\frac{\theta}{2}\right) \right|^{2\alpha+1} d\theta, \quad \delta_n = \frac{2\alpha + 1}{2n + 2\alpha + 1}, \quad \alpha > -1$$

and in the particular case  $\alpha = \beta = -\frac{1}{2}$ , we have  $d\sigma(\theta) = d\theta$  (Lebesgue measure) and  $\delta_n = 0 \quad \forall n = 1, 2, \dots$ . Now from Theorem 3.9, we deduce

$$\phi_n(z) = z^{(-1)^n} \phi_{n-2}(z), \quad \phi_0(z) \equiv 1, \quad \phi_1(z) \equiv \frac{1}{z}.$$

So,  $\phi_{2n}(z) = z^n$  and  $\phi_{2n+1}(z) = \frac{1}{z^{n+1}}$  for all  $n = 0, 1, 2, \dots$ . Since

$$\langle \phi_k(z), \phi_k(z) \rangle_{\sigma} = \|\phi_k(z)\|^2 = \int_{-\pi}^{\pi} d\theta = 2\pi, \quad k = 0, 1, 2, \dots$$

the orthonormal sequence  $\{\tilde{\phi}_n\}_{n=0}^{\infty}$  is given by

$$\tilde{\phi}_{2n}(z) = \frac{z^n}{\sqrt{2\pi}}, \quad \tilde{\phi}_{2n+1}(z) = \frac{1}{\sqrt{2\pi}z^{n+1}}, \quad n = 0, 1, 2, \dots$$

and from (3.11) we obtain:

1. If  $n$  is even:

$$R_n(z) = \alpha\phi_n(z) + \beta\phi_{n-1}(z) = \frac{\alpha z^n + \beta}{z^{n/2}} = 0 \iff \alpha z^n + \beta = 0, \quad \alpha \neq 0, \quad \left| \frac{\beta}{\alpha} \right| = 1.$$

2. If  $n$  is odd:

$$R_n(z) = \alpha\phi_n(z) + \beta\phi_{n-1}(z) = \frac{\beta z^n + \alpha}{z^{n+1}}, \quad \beta \neq 0, \quad \left| \frac{\alpha}{\beta} \right| = 1.$$

Thus, the nodes  $\{z_j\}_{j=1}^n$  of any  $n$ -point Szegő formula for the Lebesgue measure are the roots of  $z^n + \tau = 0$ ,  $|\tau| = 1$  and the weights  $\{\lambda_j\}_{j=1}^n$  are given by

$$\lambda_j = \frac{1}{\sum_{k=0}^{n-1} \left| \tilde{\phi}_k(z_j) \right|^2} = \frac{1}{\sum_{k=0}^{n-1} \left| \frac{z_j^k}{\sqrt{2\pi}} \right|^2} = \frac{2\pi}{n}, \quad j = 1, \dots, n$$

(compare with [20, pp. 73-74]).

**4. A connection with the split Levinson algorithm.** In general, when one needs to compute an  $n$ -point Szegő formula for a measure  $\sigma$  on  $\mathbb{T}$ , we start from the usual known information about  $\sigma$ , i.e., its trigonometric moments  $\mu_k = \int_{-\pi}^{\pi} e^{-ik\theta} d\sigma(\theta)$ ,  $k \in \mathbb{Z}$ , so that, as already seen through Section 3, we essentially need to calculate the (monic) Szegő polynomials  $\rho_0, \dots, \rho_n$ . As a rule, these polynomials are not explicitly known (unless for some particular measures  $\sigma$ ). Hence they should be computed from the relations (3.13). Thus, the well known Levinson algorithm arises [21]. Assume now (as will be done in the rest of the section) that the trigonometric moments  $\mu_k$  are real (a very common situation as illustrated in Section 5). In this case, Levinson's algorithm is redundant, in the sense that more operations than the required ones are carried out. To overcome this drawback a modification of this algorithm was studied giving rise to the so-called "split Levinson algorithm" (see [10]). Here, certain polynomials closely related to Szegő polynomials are computed so that the number of required operations is halved. Let us next see how these polynomials arise in the context of the construction of Szegő quadrature formulas. Indeed, in order to compute the nodes of an  $n$ -point Szegő formula and since we are dealing with real moments, it seems natural to handle polynomials (whose zeros provide us with the required nodes) with real coefficients. In other words, the parameters  $\alpha$  and  $\beta$  in (3.11) will be chosen so that

$$R_n(z) = \alpha\phi_n(z) + \beta\phi_{n-1}(z) = \frac{N_n(z)}{z^{p(n)}} = \frac{\lambda_n(\rho_n(z) + \tau_n\rho_n^*(z))}{z^{p(n)}},$$

where  $\lambda_n \neq 0$ ,  $\tau_n \in \mathbb{R}$  and  $|\tau_n| = 1$ .

Set first  $\tau_n = 1$ . Then by (3.13) it follows (now  $\delta_n \in \mathbb{R}$ )

$$N_n(z) = \lambda_n(1 + \delta_n) [\rho_{n-1}^*(z) + z\rho_{n-1}(z)].$$

Thus, taking  $\lambda_n = \frac{1}{1+\delta_n}$ , one has

$$(4.1) \quad N_n(z) = \rho_{n-1}^*(z) + z\rho_{n-1}(z) = P_n(z).$$

By [10] it can be shown that the polynomial  $P_n(z)$  represents the basic ingredient in the “symmetric version” of the Levinson-split algorithm (here it should be noted that from the context of the solution of Toeplitz systems,  $\rho_n^*(z)$  is called the “predictor polynomial”). For this reason,  $P_n(z)$  given by (4.1) is sometimes called the “first singular predictor polynomial of degree  $n$ ”, since roughly speaking it can be interpreted as though in (3.13),  $\delta_n$  is forced to take the value 1.

Setting  $P_n(z) = \sum_{j=0}^n p_{n,j} z^j$  in [10] it is proved that the sequence  $\{P_n(z)\}$  satisfies the following three-term recurrence relation,

$$(4.2) \quad P_{n+1}(z) - (1+z)P_n(z) + \alpha_n z P_{n-1}(z) = 0, \quad n \geq 1,$$

with initial conditions  $P_0(z) \equiv 2$ ,  $P_1(z) = 1+z$ , and where  $\alpha_n = \frac{\gamma_n}{\gamma_{n-1}}$  with  $\gamma_n = \sum_{i=0}^n \mu_i p_{n,i}$  for  $n \geq 1$  and  $\gamma_0 = \mu_0$ .

On the other hand, for  $\tau_n = -1$ , again by (3.13) it follows that

$$N_n(z) = \lambda_n (\delta_n - 1) [\rho_{n-1}^*(z) - z \rho_{n-1}(z)].$$

Thus, with  $\lambda_n = \frac{1}{\delta_n - 1}$ , one can write for  $n \geq 1$ ,

$$(4.3) \quad N_n(z) = \rho_{n-1}^*(z) - z \rho_{n-1}(z) = \tilde{P}_n(z).$$

Now,  $\tilde{P}_n(z)$  is the basis of the “antisymmetric version” of the Levinson-split algorithm and used to be called the “second singular predictor polynomial” (set  $\delta_n = -1$  in (3.13)). By defining  $\tilde{P}_0(z) \equiv 1$ , the polynomials  $\tilde{P}_n(z)$  for  $n = 1, 2, \dots$  are given by  $\tilde{P}_1(z) = 1 - z$ ,  $\tilde{P}_2(z) = 1 - z^2$  and for  $n \geq 2$  it holds

$$(4.4) \quad \tilde{P}_{n+1}(z) - (1+z)\tilde{P}_n(z) + \tilde{\alpha}_n z \tilde{P}_{n-1}(z) = 0,$$

where  $\tilde{\alpha}_n = \frac{\tilde{\gamma}_n}{\tilde{\gamma}_{n-1}}$  with  $\tilde{\gamma}_n = \sum_{i=0}^n \mu_i \tilde{p}_{n,i}$  and  $\tilde{P}_n(z) = \sum_{j=0}^n \tilde{p}_{n,j} z^j$ .

In order to illustrate relations (4.2) and (4.4), let us consider the Lebesgue measure, i.e.,  $d\sigma(\theta) = d\theta$ . Then,

$$\mu_k = \int_{-\pi}^{\pi} e^{-ik\theta} d\theta = 0, \quad k \geq 1, \quad \mu_0 = 2\pi.$$

From (4.2) it can be easily checked that  $P_k(0) = p_{k,0} = 1$  for  $k \geq 1$ , which gives  $\gamma_k = \mu_0$  for  $k \geq 0$ , and hence  $\alpha_k = 1$  for  $k \geq 1$ . Then (4.2) becomes

$$(4.5) \quad P_{n+1}(z) - (1+z)P_n(z) + zP_{n-1}(z) = 0, \quad n \geq 1,$$

with  $P_0(z) \equiv 2$  and  $P_1(z) \equiv 1$ . Since (4.5) is a linear second order finite difference equation with constant coefficients, it holds that

$$P_n(z) = C_1 + C_2 z^n, \quad C_1, C_2 \in \mathbb{R}.$$

From the initial conditions, one has  $C_1 = C_2 = 1$ , so that an explicit expression for the polynomial  $P_n(z)$  is obtained, namely

$$(4.6) \quad P_n(z) = z^n + 1, \quad n = 0, 1, 2, \dots$$

Similarly for the sequence  $\{\tilde{P}_n(z)\}$  one has

$$(4.7) \quad \tilde{P}_{n+1}(z) - (1+z)\tilde{P}_n(z) + z\tilde{P}_{n-1}(z) = 0, \quad n \geq 2,$$

with  $\tilde{P}_1(z) = 1 - z$  and  $\tilde{P}_2(z) = 1 - z^2$ . Thus, for  $n \geq 2$ ,

$$(4.8) \quad \tilde{P}_n(z) = 1 - z^n.$$

Clearly, (4.8) holds for  $n = 1$ .

When dealing with other measures  $\sigma$ , in general, equation (4.5) or (4.7) cannot be explicitly solved so that we need to compute recursively the polynomials  $\{P_n(z)\}_{n=0}^{\infty}$  or  $\{\tilde{P}_n(z)\}_{n=0}^{\infty}$  (for details concerning stability, see [10]).

We conclude this section rewriting the polynomials  $\{P_n(z)\}_{n=0}^{\infty}$  and  $\{\tilde{P}_n(z)\}_{n=0}^{\infty}$  in terms of the orthogonal Laurent polynomials  $\phi_n(z)$  and  $\phi_{n-1}(z)$ .

**THEOREM 4.1.** *Let  $\sigma$  be a measure on  $\mathbb{T}$  with real moments and let  $\{\phi_k\}_{k=0}^{\infty}$  be the sequence of monic orthogonal Laurent polynomials. Let  $P_n(z)$  and  $\tilde{P}_n(z)$  be the first and second singular predictor polynomials of degree  $n$  respectively. Then*

1.  $P_n(z) = z^{E[\frac{n+1}{2}]} [\phi_n(z) + (1 - \delta_n)\phi_{n-1}(z)]$
2.  $\tilde{P}_n(z) = z^{E[\frac{n+1}{2}]} (-1)^{n+1} [\phi_n(z) - (1 + \delta_n)\phi_{n-1}(z)]$

*Proof.*

1. Recall that  $\phi_{2n}(z) = \frac{1}{z^n} \rho_{2n}(z)$  and  $\phi_{2n+1}(z) = \frac{1}{z^{n+1}} \rho_{2n+1}^*(z)$ . Now for  $\alpha$  and  $\beta$  complex numbers, consider

$$R_n(z) = \alpha\phi_n(z) + \beta\phi_{n-1}(z).$$

Then, for  $n = 2m$  it follows that

$$R_{2m}(z) = \frac{\alpha\rho_{2m}(z) + \beta\rho_{2m-1}^*(z)}{z^m}$$

and for  $n = 2m + 1$

$$R_{2m+1}(z) = \frac{\alpha\rho_{2m+1}^*(z) + \beta z\rho_{2m}(z)}{z^{m+1}}.$$

Now, making use of (3.13), we have

$$R_{2m}(z) = \frac{\alpha z\rho_{2m-1}(z) + (\alpha\delta_{2m} + \beta)\rho_{2m-1}^*(z)}{z^m}.$$

Thus, taking  $\alpha = 1$  and  $\beta = 1 - \delta_{2m}$ , it follows that

$$R_{2m}(z) = \frac{P_{2m}(z)}{z^m},$$

or equivalently,

$$(4.9) \quad \begin{aligned} P_{2m}(z) &= z^m R_{2m}(z) = z^m [\alpha\phi_{2m}(z) + \beta\phi_{2m-1}(z)] = \\ &= z^m [\phi_{2m}(z) + (1 - \delta_{2m})\phi_{2m-1}(z)], \quad m \geq 1. \end{aligned}$$

Similarly for  $n = 2m + 1$ , taking  $\alpha = 1$  and  $\beta = 1 - \delta_{2m+1}$ , one can write

$$(4.10) \quad P_{2m+1}(z) = z^{m+1} [\phi_{2m+1}(z) + (1 - \delta_{2m+1})\phi_{2m}(z)].$$

So, from (4.9) and (4.10) the proof follows.

2. It can be proved in a similar way

□

**5. Numerical Examples.** Suppose that the evaluation of integrals of the form

$$(5.1) \quad \int_{-1}^1 \frac{f(x)}{(x+\lambda)^r} \omega(x) dx, \quad r = 1, 2, 3, \dots$$

where  $f(x)$  is a continuous function on  $[-1, 1]$ ,  $\omega(x) \geq 0$  a weight function on this interval and  $\lambda \in \mathbb{R}$  such that  $|\lambda| > 1$ , is required. Two methods are to be proposed:

Method 1

Let  $a$  and  $b$  be any two positive real numbers, such that

$$\frac{b}{a} = \frac{|\lambda| + 1}{|\lambda| - 1},$$

and introduce the change of variable  $x = h(t) = \frac{|\lambda|}{\lambda} \cdot \left( \frac{2t-b-a}{b-a} \right)$ . Then

$$(5.2) \quad \int_{-1}^1 \frac{f(x)}{(x+\lambda)^r} \omega(x) dx = \left( \frac{b-a}{2} \right)^r \left( \frac{|\lambda|}{\lambda} \right)^r \int_a^b g(t) \frac{\mu(t)}{t^r} dt,$$

where  $g(t) = f(h(t))$  and  $\mu(t) = \omega(h(t))$ . The integral of the right-hand side in formula (5.2) can be approximated by a Gauss-Laurent quadrature formula.

Method 2

Set  $z = e^{i\theta}$ ,  $x = \frac{1}{2}(z + z^{-1}) = \cos\theta$  and define the symmetric weight function  $\sigma(\theta)$  on  $[-\pi, \pi]$  by

$$(5.3) \quad \sigma(\theta) = \frac{2^r |\alpha|^r \omega(\cos\theta) |\sin\theta|}{|z - \alpha|^r},$$

where  $\alpha \neq 0$  is the root of the equation  $z^2 + 2\lambda z + 1 = 0$  with  $|\alpha| < 1$ , that is,

$$\alpha = \begin{cases} -\lambda + \sqrt{\lambda^2 - 1}, & \text{if } \lambda > 0, \\ -\lambda - \sqrt{\lambda^2 - 1}, & \text{if } \lambda < 0. \end{cases}$$

Then

$$(5.4) \quad \int_{-1}^1 \frac{f(x)}{(x+\lambda)^r} \omega(x) dx = \int_{-\pi}^{\pi} g(e^{i\theta}) \sigma(\theta) d\theta$$

with

$$g(e^{i\theta}) = \frac{1}{2} f \left( \frac{e^{i\theta} + e^{-i\theta}}{2} \right).$$

The integral on the right side in formula (5.4) can be approximated by an  $n$ -point Szegő quadrature formula.

For the numerical experiments we will consider two particular weight functions:

Case 1:  $\omega(x) = \frac{1}{\sqrt{1-x^2}}$ : Tchebyshev weight function of the first kind ( $x \in (-1, 1)$ ). The weight function  $\mu(t) = \omega(h(t))$  is the strong Tchebyshev weight function of the first kind on  $(a, b)$  given by

$$\mu(t) = \frac{1}{\sqrt{b-t}\sqrt{t-a}}, \quad 0 < a < b < +\infty$$

In this particular situation explicit formulas for the weights and nodes are known (see [26]):

$$x_{n+1-m} = (\beta + \alpha\nu_m) + \sqrt{(\beta + \alpha\nu_m)^2 - \beta^2}, \quad x_m = \frac{\beta^2}{x_{n+1-m}},$$

where  $m = 1, 2, \dots, p(n) = E\left[\frac{n+1}{2}\right]$ ,

$$\alpha = \frac{(\sqrt{b} - \sqrt{a})^2}{4}, \quad \beta = \sqrt{ab}, \quad \nu_m = 1 + \cos\left(\frac{2m-1}{n}\pi\right)$$

and

$$A_k = \frac{2\pi}{n} \frac{x_k}{x_k + \beta}, \quad k = 1, 2, \dots, n, \quad n \geq 1.$$

On the other hand, the weight function  $\sigma(\theta)$  on  $[-\pi, \pi]$  given by (5.3) is a rational modification of the Lebesgue measure, namely

$$\sigma(\theta) = \frac{2^r |\alpha|^r}{|z - \alpha|^r}, \quad z = e^{i\theta}.$$

Choosing the particular values  $\lambda = 1.1, 1.01$ ,  $r = 1, 2$ , the smooth function  $f(x) = e^x$  and  $n = 10$  (number of nodes) fixed we will approximate the next four integrals where exact values were computed with MATHEMATICA:

$$I_1 = \int_{-1}^1 \frac{e^x}{\sqrt{1-x^2}|x+1.1|} dx = 4.398898203 \dots$$

$$I_2 = \int_{-1}^1 \frac{e^x}{\sqrt{1-x^2}|x+1.01|} dx = 10.26398779 \dots$$

$$I_3 = \int_{-1}^1 \frac{e^x}{\sqrt{1-x^2}|x+1.1|^2} dx = 15.06117492 \dots$$

$$I_4 = \int_{-1}^1 \frac{e^x}{\sqrt{1-x^2}|x+1.01|^2} dx = 414.4873459 \dots$$

For the Tchebyshev weight function of the first kind  $\omega(x)$ , the nodes and weights of the Gaussian quadrature formula have explicit expressions. Because of the presence of a singularity near the interval of integration, we can expect to obtain poor results using this formula, as is established in the next tables:



Table 1

$r = 1$	Approximation	Exact value
$\lambda = 1.01$	9.36117219...	10.2639877...
$\lambda = 1.1$	<u>4.39825773...</u>	4.39889820...

Table 2

$r = 2$	Approximation	Exact value
$\lambda = 1.01$	247.983129...	414.487345...
$\lambda = 1.1$	<u>15.0292310...</u>	15.0611749...

In order to improve the slow convergence in this situation we use the two above mentioned methods:

Results for Method 1

Choosing  $a = 1$  implies  $b = 21$  when  $\lambda = 1.1$  and  $b = 201$  when  $\lambda = 1.01$ .

The nodes and weights obtained are:

Table 3

$\lambda = 1.1$	$n = 10$
Nodes	Weights
1.015968076	0.1140210017
1.153792215	0.1263777081
1.490163636	0.1541804078
2.180737182	0.2025926653
3.529658620	0.2733833654
5.949583872	0.3549351653
9.629771149	0.4257258654
14.09241206	0.4741381229
18.20085084	0.5019408226
20.66994082	0.5142975291

Table 4

$\lambda = 1.01$	$n = 10$
Nodes	Weights
196.7288912	0.5860819973
164.9922240	0.5786005592
113.4050389	0.5584974065
60.84072229	0.5095745959
24.34950355	0.3971049908
8.254788423	0.2312135400
3.303708313	0.1187439348
1.772408016	0.0698211241
1.218239230	0.0497179710
1.021710633	0.0422365330

The approximations of the above four integrals are:

Table 5

$r = 1$	Approximation	Exact value
$\lambda = 1.01$	<u>10.26398781..</u>	10.26398779..
$\lambda = 1.1$	<u>4.398898202..</u>	4.398898203..

Table 6

$r = 2$	Approximation	Exact value
$\lambda = 1.01$	<u>414.487344..</u>	414.4873459..
$\lambda = 1.1$	<u>15.06117499..</u>	15.06117492..

Results for Method 2

Expressions for the trigonometric moments for  $r = 1, 2$  are given by

$$\mu_k = \int_{-\pi}^{\pi} e^{-ik\theta} \sigma(\theta) d\theta = \begin{cases} \frac{4\pi|\alpha|}{1-\alpha^2} \alpha^k, & \text{if } r = 1, \\ \frac{8\pi\alpha^{k+2}}{(1-\alpha^2)^3} (k(1-\alpha^2) + 1 + \alpha^2), & \text{if } r = 2. \end{cases}$$

By using the split Levinson algorithm computed in FORTRAN with double precision, the following nodes and weights are obtained:

Table 7

$r = 1, \lambda = 1.1$	$n = 10$
Nodes	Weights
$0.5231956081 \pm 0.8522126235i$	0.4170158485
$-0.9718655888 \pm 0.2355361486i$	3.8990461817
$-0.6941445801 \pm 0.7198356076i$	1.5283999088
$0.9425118124 \pm 0.3341728347i$	0.3334871136
$-0.1205684668 \pm 0.992705014i$	0.6775681574

Table 8

$r = 1, \lambda = 1.01$	$n = 10$
Nodes	Weights
$0.5077320439 \pm 0.8615150443i$	0.4552583228
$-0.1556028488 \pm 0.9878196968i$	0.8023120227
$0.9406267445 \pm 0.3394426719i$	0.3550340683
$-0.7396737275 \pm 0.672965658i$	2.4403459272
$-0.9871949772 \pm 0.1595182654i$	18.1061357088

Table 9

$r = 2, \lambda = 1.1$	$n = 10$
Nodes	Weights
$-0.7826063935 \pm 0.6225168535i$	5.7286222677
$-0.2523182279 \pm 0.9676443106i$	0.9628770791
$0.9315406380 \pm 0.3636372365i$	0.1801413420
$0.4439464841 \pm 0.8960533016i$	0.3067180364
$-0.9823049310 \pm 0.1872886076i$	28.7314933198

Table 10

$r = 2, \lambda = 1.01$	$n = 10$
Nodes	Weights
$0.9264937183 \pm 0.376310231i$ $-0.8669062623 \pm 0.4984711951i$ $0.4037996706 \pm 0.9148474332i$ $-0.3361546099 \pm 0.9418068158i$ $-0.9954580194 \pm 0.0952015315i$	0.2056746266 30.7425691490 0.3833193864 1.6432603190 1080.4916890184

The approximation of the above four integrals are

Table 11

$r = 1$	Approximation	Exact value
$\lambda = 1.01$	<u>10.26398785..</u>	10.26398779..
$\lambda = 1.1$	<u>4.398898196..</u>	4.398898203..

Table 12

$r = 2$	Approximation	Exact value
$\lambda = 1.01$	<u>414.4873471..</u>	414.4873459..
$\lambda = 1.1$	<u>15.06117499..</u>	15.06117492..

From Tables 5-6 and 11-12 one can see that both methods provide us with similar numerical results. In both cases results given by Gaussian formulas are strongly improved.

Case 2:  $\omega(x) = \sqrt{1-x^2}$ : Tchebyshev weight function of the second kind ( $x \in [-1, 1]$ ). The weight function  $\mu(t) = \omega(h(t))$  is the strong Tchebyshev weight function of the second kind on  $[a, b]$  given by

$$\mu(t) = \sqrt{b-t}\sqrt{t-a}.$$

Choosing the particular values  $\lambda = 1.1, 1.01, r = 1$ , the smooth function  $f(x) = e^x$  and again  $n = 10$  (number of nodes) fixed, we will approximate the next two integrals with exact values computed with MATHEMATICA:

$$I_1 = \int_{-1}^1 \frac{e^x \sqrt{1-x^2}}{|x+1.1|} dx = 1.67594127475 \dots$$

$$I_2 = \int_{-1}^1 \frac{e^x \sqrt{1-x^2}}{|x+1.01|} dx = 2.03543204817 \dots$$

For the Tchebyshev weight function of the second kind  $\omega(x)$ , the nodes and weights of the Gaussian quadrature formula have also explicit expressions. As in case 1, because of the presence of a singularity near the interval of integration, we can expect to obtain poor results using this formula as displayed in the next table:

Table 13

$r = 1$	Approxiamtion	Exact value
$\lambda = 1.01$	<u>2.421155521</u> ...	2.03543204817 ...
$\lambda = 1.1$	<u>1.608794159</u> ...	1.67594127475 ...

Again, in order to improve the slow convergence in this situation we use the two mentioned methods:

Results for Method 1

From the relation

$$\int_{-1}^1 \frac{f(x)\sqrt{1-x^2}}{|x+\lambda|^r} dx = \int_{-1}^1 \frac{g(x)}{\sqrt{1-x^2}|x+\lambda|^r} dx$$

with

$$g(x) = f(x)(1-x^2),$$

the mentioned explicit formulas for the nodes and weights in case 1 can be used. The approximation of the above two integrals are

Table 14

$r = 1$	Approximation	Exact value
$\lambda = 1.01$	<u>2.035432052</u> ...	2.03543204817 ...
$\lambda = 1.1$	<u>1.675941276</u> ...	1.67594127475 ...

Results for Method 2

Expressions for the trigonometric moments are given by

$$\mu_k = \begin{cases} 2\pi|\alpha|, & \text{if } k = 0, \\ \pi\alpha|\alpha|, & \text{if } k = 1, \\ -\pi|\alpha|\alpha^{k-2}(1-\alpha^2), & \text{if } k \geq 2, \end{cases}$$

when  $r = 1$ , and

$$\mu_k = \begin{cases} \frac{-4\pi}{1-\alpha^2}, & \text{if } k = 0, \\ -2\pi\frac{\alpha^k}{1-\alpha^2} (2\alpha + k - 1 - \alpha^2(k+3)), & \text{if } k \geq 1, \end{cases}$$

when  $r = 2$ . By the split Levinson algorithm computed in FORTRAN with double precision, nodes and weights are given by:

Table 15

$r = 1, \lambda = 1.1$	$n = 10$
Nodes	Weights
$0.8473567133 + \pm 0.5310241053i$	0.0809799064
$-0.9095350917 + \pm 0.4156271369i$	0.4250829076
$-0.5968012792 + \pm 0.8023890784i$	0.6749529041
$0.4377179772 + \pm 0.8991123247i$	0.2923674821
$-0.0996095352 + \pm 0.9950266029i$	0.5427101048

and,

Table 16

$r = 1, \lambda = 1.01$	$n = 10$
Nodes	Weights
$0.8432691294 + \pm 0.5374915584i$	0.0884264387
$-0.9380253965 + \pm 0.3465665239i$	0.8084274344
$-0.6342873464 + \pm 0.7730973821i$	0.8785194976
$0.4228259917 + \pm 0.906210892i$	0.3244615048
$-0.1278951430 + \pm 0.9917876952i$	0.6277760750

Finally the approximation of the two integrals are displayed in:

Table 17

$r = 1$	Approximation	Exact value
$\lambda = 1.01$	<u>2.03543204774</u> ...	2.03543204817...
$\lambda = 1.1$	<u>1.67594127382</u> ...	1.67594127475...

Again, numerical results offered by both methods are quite similar.

**Acknowledgements.** The authors thank the referee for valuable suggestions which have contributed to improve the final form of this paper. The work of this author was partially supported by the Scientific Research Project BFM2001-3411 of the Spanish Ministry of Science and Technology.

REFERENCES

- [1] N.I. AKHIEZER, *The classical moment problem and some related questions in analysis*, Hafner, New York, 1965.
- [2] A. BULTHEEL, C. DÍAZ-MENDOZA, P. GONZÁLEZ-VERA AND R. ORIVE, *On the convergence of certain Gauss-type quadrature formulas for unbounded intervals*, Math. Comp., 69 (2000), pp. 721-747.
- [3] A. BULTHEEL, P. GONZÁLEZ-VERA, E. HENDRIKSEN AND O. NJÅSTAD, *Orthogonal rational functions and quadrature on the real half-line*, Journal of Complexity, to appear.

- [4] A. BULTHEEL, C. DÍAZ-MENDOZA, P. GONZÁLEZ-VERA AND R. ORIVE, *Quadrature on the half-line and two point Padé approximants to Stieljes functions, Part II: Convergence*, J. Comput. Appl. Math., 77 (1997), pp. 53-76.
- [5] A. BULTHEEL, C. DÍAZ-MENDOZA, P. GONZÁLEZ-VERA AND R. ORIVE, *Orthogonal Laurent polynomials and quadrature formulas for unbounded intervals: I. Gauss-type formulas*, Rocky Mountain J. Mathematics, to appear.
- [6] A. BULTHEEL, P. GONZÁLEZ-VERA AND R. ORIVE, *Quadrature on the half-line and two point Padé approximants to Stieljes functions, Part I: Algebraic aspects*, J. Comput. Appl. Math., 65 (1996), pp. 57-72.
- [7] L. COCHRAN AND S. CLEMENT COOPER, *Orthogonal Laurent polynomials on the real line*, in S. Clement Cooper, and W.J. Thron, eds., Lecture Notes in Pure and Applied Mathematics., vol. 154, Marcel Dekker, 1994, pp. 47-100.
- [8] L. DARUIS, P. GONZÁLEZ-VERA AND O. NJÅSTAD., *Szegő quadrature formulas for certain Jacobi-type weight functions*, Mathematics of Computation, 71 (2001), No. 238, pp. 683-701.
- [9] L. DARUIS AND P. GONZÁLEZ-VERA, *Interpolatory quadrature formulas on the unit circle for Chebyshev weight functions*, Numer. Math., 90 (2002), pp. 641-664.
- [10] P. DELSARTE, Y. GENIN, *The Split Levinson Algorithm*, IEEE Trans Acoust. Speech Signal Proc., 34 (1986), pp. 470-478.
- [11] W. GAUTSCHI, *A survey of Gauss-Christoffel quadrature formulae*, in Butzer, P.L., Fehér, F., eds., E.B. Christoffel - The influence of his work in mathematics and the physical sciences, Birkhäuser Verlag, Basel, 1981, pp. 72-147.
- [12] W. GAUTSCHI, *Gauss-type quadrature rules for rational functions*, in H. Brass, G. Hämmerlin, eds., Numerical Integration IV, International Series of Numerical Mathematics, vol. 112, Birkhäuser, Basel, 1993, pp. 111-130.
- [13] P. GONZÁLEZ-VERA, O. NJÅSTAD AND J.C. SANTOS-LEÓN, *Some results about numerical quadrature on the unit circle*, Advances in Computational Mathematics, 5 (1996), pp. 297-328.
- [14] W.B. GRAGG, *Positive definite Toeplitz matrices, the Arnoldi process for isometric operators, and Gaussian quadrature on the Unit Circle*, J. Comput. Appl. Math., 46 (1993), pp. 183-198.
- [15] P.E. GUSTAFSON AND B.A. HAGLER, *Gaussian quadrature rules and numerical examples for strong extensions of mass distribution functions*, J. Comput. Appl. Math., 105 (1999), pp. 317-326.
- [16] W.B. JONES, O. NJÅSTAD AND W.J. THRON, *Two-point Padé expansions for a family of analytic functions*, J. Comput. Appl. Math., 9 (1983), pp. 105-123.
- [17] W.B. JONES, O. NJÅSTAD AND W.J. THRON, *Moment theory, orthogonal polynomials, quadrature, and continued fractions associated with the unit circle*, Bull. London Math. Soc., 21 (1989), pp. 113-152.
- [18] W.B. JONES, W.J. THRON AND H. WADELAND, *A strong Stieljes moment problem*, Trans. of the AMS, 261 (1980), pp. 503-528.
- [19] W.B. JONES AND W.J. THRON, *Orthogonal Laurent polynomials and Gaussian quadrature*, in Quantum Mechanics in Mathematics, Chemistry and Physics, K.E. Gustafson et al., eds., Plenum Publ. Comp., 1981, pp. 449-455.
- [20] V.I. KRYLOV, *Approximate Calculation of Integrals*, The MacMillan Company, New York, 1962.
- [21] N. LEVINSON, *The Wiener RMS (root mean square) error criterion in filter design and prediction*, J.Math. Phys., 25 (1947), pp. 261-278.
- [22] G. LÓPEZ-LAGOMASINO AND A. MARTÍNEZ-FINKELSHEIN, *Rate of convergence of two-point Padé approximants and logarithmic asymptotics of Laurent-type orthogonal polynomials*, Constr. Approx., 11 (1995), pp. 255-286.
- [23] G. LÓPEZ-LAGOMASINO AND J. ILLÁN-GONZÁLEZ, *Quadrature formulas for unbounded intervals*, Rev. Cienc. Mat., 3 (1982), pp. 29-47.
- [24] O. NJÅSTAD AND W.J. THRON, *The theory of sequences of orthogonal L-polynomials*, in Padé approximants and Continued Fractions, H. Waadeland et al., eds., Det Kongelige Norske Videnskabers Selskab Skrifter, No. 1, 1983, pp. 54-91.
- [25] J.C. SANTOS, *Computation of integrals over the unit circle with nearby poles*, Appl. Numer. Math., 36 (2001), No. 2-3, pp. 179-195.
- [26] A. SRI RANGA, *Another quadrature rule of highest algebraic degree of precision*, Numer. Math., 68 (1991), pp. 283-294.
- [27] A. SRI RANGA, *J-Fractions and strong moment problems*, in W.J. Thron, ed., Analytic Theory of Continued Fractions, II, Proceedings Pitlochry and Aviemore 1985. Lecture Notes in Mathematics Springer-Verlag, No. 1199, 1986, pp. 269-284.
- [28] G. SZEGÖ, *Orthogonal polynomials*, A.M.S., 4th ed., vol. 23, Colloquium Publications, 1975.
- [29] G. SZEGÖ, *On biorthogonal systems of trigonometric polynomials*, Magyar. Tud. Akad. Kutató Int. Közl., 8 (1963), pp. 255-273.
- [30] W.J. THRON, *L-polynomials orthogonal on the unit circle*, in Nonlinear Methods and Rational Approximation, A. Cuyt, ed., D. Reidel Publishing Company, 1988, pp. 271-278.