

PROOF OF A CONJECTURE OF CHAN, ROBBINS, AND YUEN *

DORON ZEILBERGER †

Abstract. Using the celebrated Morris Constant Term Identity, we deduce a recent conjecture of Chan, Robbins, and Yuen (math.CO/9810154), that asserts that the volume of a certain $n(n - 1)/2$ -dimensional polytope is given in terms of the product of the first $n - 1$ Catalan numbers.

Key words. combinatorics, Catalan numbers, polytope.

AMS subject classifications. 05-XX, 52B05.

1. Main Result. Chan, Robbins, and Yuen [1] conjectured that the cardinality of a certain set of triangular arrays \mathcal{A}_n defined in pp. 6-7 of [1] equals the product of the first $n - 1$ Catalan numbers. It is easy to see that their conjecture is equivalent to the following *constant term identity* (for any rational function $f(z)$ of a variable z , $CT_z f(z)$ is the coeff. of z^0 in the formal Laurent expansion of $f(z)$ (that always exists)):

$$(1.1) \quad CT_{x_n} \dots CT_{x_1} \prod_{i=1}^n (1 - x_i)^{-2} \prod_{1 \leq i < j \leq n} (x_j - x_i)^{-1} = \prod_{i=1}^n \frac{1}{i+1} \binom{2i}{i}.$$

But this is just the special case $a = 2, b = 0, c = 1/2$, of the *Morris Identity* [2] (where we made some trivial changes of discrete variables, and ‘shadowed’ it)

$$(1.2) \quad CT_{x_n} \dots CT_{x_1} \prod_{i=1}^n (1 - x_i)^{-a} \prod_{i=1}^n x_i^{-b} \prod_{1 \leq i < j \leq n} (x_j - x_i)^{-2c} = \frac{1}{n!} \prod_{j=0}^{n-1} \frac{\Gamma(a + b + (n - 1 + j)c)\Gamma(c)}{\Gamma(a + jc)\Gamma(c + jc)\Gamma(b + jc + 1)}.$$

To show that the right side of (1.2) reduces to the right side of (1.1) upon the specialization $a = 2, b = 0, c = 1/2$, do the plugging in the former and call it M_n . Then manipulate the products to simplify M_n/M_{n-1} , and then use *Legendre’s duplication formula* $\Gamma(z)\Gamma(z + 1/2) = \Gamma(2z)\Gamma(1/2)/2^{2z-1}$ three times, and *voilà*, up pops the Catalan number $\binom{2n}{n}/(n + 1)$. □

REMARK 1.1. *By converting the left side of (1.2) into a contour integral, we get the same integrand as in the Selberg integral (with $a \rightarrow -a, b \rightarrow -b - 1, c \rightarrow -c$). Aomoto’s proof of the Selberg integral (SIAM J. Math. Anal. **18**(1987), 545-549) goes verbatim.*

REMARK 1.2. *Conjecture 2 in [1] follows in the same way, from (the obvious contour-integral analog of) Aomoto’s extension of Selberg’s integral. Introduce a new variable t , stick $CT_t t^{-k}$ in front of (1.1), and replace $(1 - x_i)^{-2}$ by $(1 - x_i)^{-1}(t + x_i/(1 - x_i))$.*

REMARK 1.3. *Conjecture 3 follows in the same way from another specialization of (1.2).*

†Department of Mathematics, Temple University, Philadelphia, PA 19122, USA. (zeilberg@math.temple.edu). <http://www.math.temple.edu/~zeilberg/> . Supported in part by the NSF.

*Received November 1, 1998. Accepted for publication December 1, 1999. Recommended by F. Marcellán.

REFERENCES

- [1] CLARA S. CHAN, DAVID P. ROBBINS, AND DAVID S. YUEN, *On the volume of a certain polytope*, math.CO/9810154.
- [2] WALTER MORRIS, “*Constant term identities for finite and affine root systems, conjectures and theorems*”, Ph.D. thesis, University of Wisconsin, Madison, Wisconsin, 1982.